

Steps Toward a Common Near-Infrared Photometric System

A. T. Tokunaga

*Institute for Astronomy, University of Hawaii, 2680 Woodlawn Dr.,
Honolulu, HI 96822*

W. D. Vacca

*Stratospheric Observatory for Infrared Astronomy/Universities Space
Research Association, NASA Ames Research Center, MS N211-3,
Moffett Field, CA 94035-1000*

Abstract. The proliferation of near-infrared (1–5 μm) photometric systems over the last 30 years has made the comparison of photometric results difficult. In an effort to standardize infrared filters in use, the Mauna Kea Observatories near-infrared filter set has been promoted among instrument groups through combined filter production runs. The characteristics of this filter set are summarized, and some aspects of the filter wavelength definitions, the flux density for zero magnitude, atmospheric extinction coefficients, and color correction to above the atmosphere are discussed.

1. Introduction

The foundations of modern infrared photometry at near-infrared wavelengths (1–5 μm) were built upon the pioneering work of Johnson and his collaborators (Johnson 1966; Johnson et al. 1966; Johnson, MacArthur, & Mitchell 1968). The near-infrared filters used in these early works were very broad, and the atmospheric absorption bands defined the effective widths of the filters. The filter profiles used by Johnson (1965) are shown in Figure 1, which demonstrates that the deep atmospheric absorption bands (mostly due to water vapor) play a significant role in determining the effective transmission of the atmosphere + telescope + filter observational system.

The problem with allowing the atmosphere to define the passband is that the atmosphere varies substantially from place to place and, at any given location, from night to night (even from hour to hour). Hence, the reproducibility of photometric observations and conversions to other photometric systems becomes difficult or impossible to achieve. To prevent these problems, the photometric bands should ideally be defined by only the filter transmission profiles (if possible). Ideally, the filter bandpasses should avoid deep telluric absorption features and any part of the telluric spectrum that has a large first derivative and/or is highly variable. In principle, filter properties can be controlled by the manufacturer, and so the specifications and characteristics of the filters are usually the

only aspect of the observational system that can remain constant from instrument to instrument and site to site. If such a set of filters were easily replicable and distributed to a large number of observatories, the comparison of photometric results from various instruments, telescopes, and sites would be much more straightforward.

There has never been an agreed upon filter set for the ground-based near-infrared filters. As a result, many types of filters are currently used, as illustrated in Figure 2, and summaries of infrared photometric systems in use are given by Bessell (2005), Glass (1999), and Tokunaga (2000). (The former two references also provide excellent introductions to infrared photometry.) As a result of this proliferation of filter sets, the comparison of results in different systems can be quite complicated. Careful attention to the details of the photometric system employed such as that provided by Moro & Munari (2000) and Fiorucci & Munari (2003) is needed. Furthermore, the labels for the filters in different sets are often the same (e.g., J , H , and K), and this can lead the unwary to mistakenly combine or compare results inappropriately. Very different filters may have been used for observations despite the common label, and corrections for the differing filter (isophotal) wavelengths must be incorporated first before comparisons can be done. For objects with very deep absorption bands or strong emission lines, direct comparison may not be possible.

The basic difficulty of having different filters lies in the intercomparison of results from the various photometric systems. In order to make accurate comparisons, all of the photometric points must be either (1) plotted at the filter wavelengths of the various systems, or (2) corrected to the same wavelength. The latter is the purpose behind color transformations. Intercomparison of results among the different systems requires color transformations such as those provided by Carpenter (2001), Hawarden et al. (2001), Stephens & Leggett (2004), and Leggett et al. (2006).

The large number of color transformations highlights the need for standardization of infrared filters. As stated in the goals for this workshop: “The astronomical community witnesses an explosive non-linear growth of observational capability ... This quantum leap and its associated uncontrolled production of diverse systems of measurement disrupts the consistency of calibrated measurements ...”. Figure 2 serves as a fitting testimony to the relevance of these statements.

Although color transformations are very helpful in combining data from different systems, the systematic effects and limited color range of the transformations make it much less desirable than having observations with a single filter set. The adoption of a standard filter set would minimize confusion about which filter set is being used and the magnitude of the color transformations. However, use of a single filter set will not eliminate the need for color transformations due to optical efficiency differences among instruments and because completely identical filters cannot be fabricated for all photometric systems.

A relatively new set of filters, the Mauna Kea Observatories near-infrared (MKO-NIR) filter set, was designed for minimizing the atmospheric absorption effects. The filter profiles are shown in Figure 1. These filters are located near the center of the high atmospheric transmission regions, and therefore the filter central wavelengths are defined by these regions.

We present here a brief summary of the properties of the MKO-NIR filter set. This filter set is gaining acceptance in the community and thus could serve as a unified near-infrared filter set in the future. We also discuss aspects of infrared photometry that may be helpful to newcomers to infrared photometry, and we discuss a method to establish an improved photometric system for the MKO-NIR filter set.

2. The MKO-NIR Filter Set

A filter set for 1–5 μm was developed initially for the Subaru and Gemini telescopes. The central wavelength and bandpass were optimized by maximizing throughput while minimizing the effects of the atmospheric transmission and emission. The method to determine this was presented by Simons & Tokunaga (2002). The 1996–2002 period was an opportune time to convince many other groups to use these filters, since there were a large number of new infrared instruments under development at the time. There was a strong motivation to combine orders for production, thus sharing the engineering and fabrication costs and thus greatly reducing the cost per filter. In addition, the filters were optimized at both 2 and 4 km altitude, and thus many observatories could use these filters. As a result these filters are now widely used at a large number of major observatories and are being employed in near-infrared sky surveys such as the UKIRT Infrared Deep Sky Survey (UKIDSS).

The detailed specifications for the fabrication of the filters and atmospheric extinction coefficients are described by Tokunaga, Simons, & Vacca (2002), while the isophotal wavelengths (for a Vega source spectrum) and absolute calibration are described by Tokunaga & Vacca (2005).

2.1. Definition of the Filter Wavelengths

There is no unique definition of the “effective wavelength” of a broad-band filter. See Golay (1974) and Tokunaga & Vacca (2005) for detailed discussions of the various filter definitions. The fundamental problem is that the filters are broad relative to the wavelength scale of the variations in the source spectra. In contrast a single pixel in a spectrograph is like a filter with a very well defined wavelength simply because its wavelength span is small compared with the variations in the spectra in most sources.

As we show below (see also Golay 1974), the isophotal wavelength is the most appropriate filter wavelength definition to use. This definition is *not* independent of the source spectrum, which presents problems, and is the reason others have used different filter wavelength definitions in an attempt to approximate the isophotal wavelength without having to incorporate the source spectrum.

To derive an expression for the isophotal wavelength we first consider the number of photo-electrons detected per second from a source with an intrinsic spectral energy distribution $F_\lambda(\lambda)$. This is given by

$$N_p = \int F_\lambda(\lambda) S(\lambda) / h\nu d\lambda \quad (1)$$

$$= \frac{1}{hc} \int \lambda F_\lambda(\lambda) S(\lambda) d\lambda \quad , \quad (2)$$

where $S(\lambda)$ is the total system response given by

$$S(\lambda) = T(\lambda)Q(\lambda)R(\lambda)A_{\text{tel}} \quad . \quad (3)$$

Here $T(\lambda)$ is the atmospheric transmission, $Q(\lambda)$ is the product of the throughput of the telescope, instrument, and quantum efficiency of the detector, $R(\lambda)$ is the filter response function, and A_{tel} is the telescope collecting area. The system response $S(\lambda)$ is equal to the relative spectral response (RSR) defined by Cohen et al. (2003a). Care is needed when taking filter response functions from various sources. For example, the 2MASS filter response functions have already been corrected for the extra factor of λ in the integral (Cohen, Wheaton, & Megeath 2003b).

If $F_\lambda(\lambda)$ and $S(\lambda)$ are both continuous, and $S(\lambda)$ is not negative over the wavelength interval, then from equation (2) and the mean value theorem for integration there exists a wavelength, λ_{iso} , such that

$$F_\lambda(\lambda_{\text{iso}}) \int \lambda S(\lambda) d\lambda = \int \lambda F_\lambda(\lambda) S(\lambda) d\lambda \quad . \quad (4)$$

Rearranging this, we obtain

$$F_\lambda(\lambda_{\text{iso}}) = \langle F_\lambda \rangle = \frac{\int \lambda F_\lambda(\lambda) S(\lambda) d\lambda}{\int \lambda S(\lambda) d\lambda} \quad , \quad (5)$$

where λ_{iso} is known as the “isophotal wavelength” and $\langle F_\lambda \rangle$ denotes the mean value of the intrinsic flux above the atmosphere (in units of $\text{W m}^{-2} \mu\text{m}^{-1}$) over the wavelength interval of the filter.

In a similar fashion,

$$F_\nu(\nu_{\text{iso}}) = \langle F_\nu \rangle = \frac{\int F_\nu(\nu) S(\nu) / \nu \, d\nu}{\int S(\nu) / \nu \, d\nu} \quad , \quad (6)$$

where ν_{iso} denotes the “isophotal frequency” and $\langle F_\nu \rangle$ denotes the mean value of the intrinsic flux above the atmosphere (in units $\text{W m}^{-2} \text{Hz}^{-1}$) over the frequency interval of the filter.

The isophotal wavelengths and frequencies, λ_{iso} and ν_{iso} , respectively, for the MKO-NIR filters are given in Table 1 (see also Tokunaga & Vacca 2005). These were computed using the model atmosphere for Vega recommended by Bohlin & Gilliland (2004) and therefore strictly apply only to A0V stars. For any other source spectrum, the isophotal wavelengths would differ from those shown in Table 1.

The isophotal wavelength is important and useful because it is the wavelength at which the monochromatic flux F_λ equals the mean flux in the passband. As stated succinctly by Golay (1974, p. 41): “The wavelength of the monochromatic intensity deduced from a heterochromatic measurement ... is ... λ_{iso} .” The use of isophotal wavelengths determined from a model of Vega provides consistency with the extensive series of papers on infrared calibration by Cohen

and his collaborators. Other definitions for the filter wavelength are discussed by Golay (1974) and Tokunaga & Vacca (2005).

We note that the UKIDSS program defined filters for Z and Y with effective wavelengths at $0.88\ \mu\text{m}$ and $1.03\ \mu\text{m}$ respectively, (Hewett et al. 2006). It is typically the case that the longest filter for CCD work is named “ Z ”. The “ Y ” filter was introduced by Hillenbrand et al. (2002) to take advantage of the atmospheric window between atmospheric absorption at 0.955 and $1.112\ \mu\text{m}$. The definition of the effective wavelength adopted by the Sloan Digital Sky Survey as well as the UKIDSS (Hewett et al. 2006) is given by (Fukugita et al. 1996, see their equation 3) and is not that of the isophotal wavelength for an A0V star. Therefore, the flux densities in Hewett et al.’s Table 7, which correspond to the flux densities of an object of zero magnitude of constant flux density over the passband, are not the monochromatic flux densities at the effective wavelengths given in their table. Thus, while the flux density zero points agree well with those given in Table 1, direct comparison of the wavelengths in Table 1 with that of Hewett et al.’s Table 7 is not appropriate.

2.2. Correction for Spectral Energy Distribution

Photometric data points are often converted to flux densities and interpreted as measurements of the monochromatic flux at wavelengths given by the “effective wavelengths” of the filters, such as the isophotal wavelengths. However, what one has measured is the “in-band” flux passing through the filter. Since photometry is unavoidably a heterochromatic (multiwavelength) measurement, observations of sources with very different spectral energy distributions cannot be directly compared at a single wavelength unless a correction is made for the different spectral energy distributions. This is clear from the definition of the isophotal wavelength, where the spectral energy distribution must be known to compute it.

The dependence of the isophotal wavelength on the source spectrum is the fundamental reason for the difficulties encountered in the interpretation and intercomparison of photometric results. To accurately know the wavelength at which a photometric data point applies, one needs to know the source spectrum, which is usually what one is attempting to determine in the first place. The application of a correction factor during photometric reductions is a means of attempting to account for the sensitivity of λ_{iso} to the slope of the source spectrum, and thereby estimating the monochromatic source flux density at the isophotal wavelength of an A0V star.

To visualize this, consider the observation of a very cool object, say $T \sim 500\ \text{K}$. Suppose the standard star is much hotter, $T \sim 10^4\ \text{K}$. Then the isophotal wavelength of the standard is inappropriate for the cool object since the flux density is weighted to the longer wavelengths compared to the standard star. If a correction is not made for the spectral energy distribution, then one will overestimate the flux density of the cool object at the isophotal wavelength of the standard. To obtain the proper flux density of the cool object at the isophotal wavelength of the standard one must apply a correction factor that reduces the flux density of the cool object. These correction factors are required for any definition of the filter wavelength, including those that are independent of the

source spectrum. Discussions of computing the correction factor can be found in Low & Rieke (1974), Hanner et al. (1984), and Glass (1999).

2.3. Conversion of Magnitudes to Flux Density

All photometry is relative, and so it is necessary to define the photometric zero point by using a star or group of stars to define zero magnitude and zero colors. In establishing the *UBV* system, Johnson & Morgan (1953) tied the *V* magnitude to nine stars of the International Photovisual system and the *U* – *B* and *B* – *V* colors to an average of six A0V stars. Most optical photometric systems assume a visual magnitude of +0.03 for Vega. However most infrared photometric systems, such as Elias et al. (1982) for the C.I.T. system and Cohen et al. (1992) for his absolute calibration work, define Vega to have zero magnitude and zero color at 1–5 μm . More recently Price et al. (2004) have chosen to tie the MSX photometry to Sirius since Vega has dust emission that affects its brightness at wavelengths longer than 12 μm . Therefore, in comparing near-infrared photometric systems, care is required to make sure that the assumptions about the visual and infrared magnitude of Vega are consistent. Cohen et al. (1992) assumed 0.0 mag for Vega in the near-infrared, but in (Cohen et al. 2003a) the nonzero magnitude of Vega at *V* was taken into account in establishing the 0.0 mag flux densities (see their Table 3).

Vega is the most well-studied star for the determination of the absolute flux density. A summary of the direct comparison of blackbody furnaces to Vega is given by Mégessier (1995), while Cohen et al. (1992) used a model atmosphere to extrapolate from the absolute calibration at *V* to the near infrared. Tokunaga & Vacca (2005) show that these two methods give the identical flux density in the near-infrared.

The flux density for Vega shown in Table 1 is from Tokunaga & Vacca (2005), and it is normalized to be consistent with the flux density of Vega determined by Cohen et al. (1992). Tokunaga & Vacca (2005) have shown that these values are consistent with the model-independent absolute calibration measurements summarized by Mégessier (1995). In addition, the absolute calibration of Cohen et al. (1992) was found to be consistent with the extensive absolute calibration conducted by the *Midcourse Space Experiment* (Price et al. 2004; Price 2004) to within 1%. Therefore the values shown in Table 1 are consistent with the extensive absolute calibration work of Cohen and his collaborators, of which Cohen, Wheaton, & Megeath (2003b) is the latest in the series.

Nevertheless, the use of a model atmosphere for Vega is fraught with difficulties because Vega is a rapid rotator and is observed nearly pole-on (Bohlin & Gilliland 2004; Peterson et al. 2006). Consequently there is a temperature gradient across the surface of Vega as viewed from Earth. Therefore, a model atmosphere with a single fixed temperature cannot provide an accurate representation of the observed flux density of Vega from the ultraviolet to the near-infrared. In fact, there is no self-consistent model of Vega that is applicable from the optical to the infrared. Nonetheless, the single temperature model assumed by Cohen et al. (1992) provides a close approximation to the near-infrared spectral energy distribution that we observe and has been validated by ground-based and space-based absolute calibration experiments. This does not imply the model atmosphere is a valid one but rather that the model atmosphere

is empirically a good match to the near-infrared spectral energy distribution of Vega.

2.4. Linearity and Extinction

In establishing the *UBV* system Johnson & Morgan (1953) employed the technique of linearly extrapolating magnitudes to above the atmosphere. A strictly linear extrapolation is not valid due to the Forbes effect; the atmospheric extinction curve is nonlinear between zero and unity airmass. This is shown explicitly for the MKO-NIR filters in Figure 3. The nonlinearity of the extinction in the near-infrared passbands was discussed by (Manduca & Bell 1979, see their Fig. 2), Young (1974), and Young, Milone, & Stagg (1994) and arises from the very strong absorption bands of water vapor in the near-infrared. As a result, all commonly used near-infrared photometric systems (Glass, Elias, Persson, UKIRT) employ the method of reducing to an airmass of 1.0. This is acceptable since all photometry is done differentially.

The effects of water vapor absorption are not completely eliminated by the design of the MKO-NIR filters. Tokunaga, Simons, & Vacca (2002) computed the extinction for the MKO-NIR filters as a function of telluric precipitable water. The atmospheric extinction coefficients is shown in Table 2 for 2 mm of precipitable water. For J , H , and K these values are smaller than for those given by Krisciunas et al. (1987) who employed older filters with wider passbands. See also the UKIRT Web site, <http://www.jach.hawaii.edu/UKIRT/astronomy/utills/exts.html>). For L' and M' the calculated extinction coefficients are identical to that observed within the errors.

Computations of the extrapolation to zero airmass as a function of water vapor from 1 to 5 mm precipitable water suggests the variations of the magnitude at zero airmass is in the range of 0.003–0.012 mag for J , H , K , K' , K_s , and L' , and in the range of 0.096–0.123 for M' for the MKO-NIR filters (Tokunaga & Vacca 2005). For nearly all work the uncertainties in the exo-atmospheric magnitudes for J , H , K , and L' are acceptable. Since the uncertainties at M' are larger, care is needed if accurate exo-atmospheric magnitudes are needed.

A complication arises for very cool stars that have deep water and other strong molecular absorption bands. In this case, there will be a color term in the extinction curve. Good discussions on measuring the color term are given by Hardie (1962) and Henden & Kaitchuck (1982), and the formal derivation of the color term in photometry is given by Golay (1974). As this color term is rarely measured, high-precision photometry of cool objects is subject to uncertainties arising from neglecting it. The importance of this color term is that when one extrapolates the observations to zero airmass (i.e., above the atmosphere), the result should be identical to observations done completely in the absence of an atmosphere.

Milone & Young (2005) have proposed relatively narrowband filters that minimize color transformations between observatories and instruments. The extrapolation to zero airmass for these filters is approximately linear. This filter set is important for observations where one wants to determine the absolute calibration to above the atmosphere. The absolute calibration work by Blackwell et al. (1983) and Selby et al. (1983) used narrowband filters exactly for this reason.

3. Setting the Photometric Zero Point for the MKO-NIR Filters

The papers by Hawarden et al. (2001), Leggett et al. (2003), and Leggett et al. (2006) give a set of standard stars measured in the MKO-NIR filter set. The magnitudes of these standard stars were largely based on the system established by Elias et al. (1982). Although the zero point of the system was not determined explicitly, it should be close to that of Elias, namely that the colors of Vega are 0.0 mag ($V - J = J - H = H - K = K - L' = L' - M' = 0.0$). We discuss in this section an approach to better establish the photometric zero point of the MKO-NIR standards relative to a set of A0V stars.

To set the zero point of the MKO-NIR photometric system, one could observe a number of A0V stars and either define Sirius to have zero colors as suggested by Cohen et al. (1992) and Price et al. (2004) or adopt an average of A0V stars to have zero colors after removing the interstellar extinction. The latter method was used by Johnson & Morgan (1953) when they defined the *UBV* system. Thus Vega has a magnitude of +0.03 mag in the Johnson system, although Bohlin & Gilliland (2004) recently determined $V = +0.026 \pm 0.008$ mag. In the former method, Sirius is proposed instead of Vega because there is no definitive model atmosphere for Vega and because Sirius is accessible to the Northern and Southern hemispheres. However Sirius and Vega are too bright to be observed with current instrumentation.

Hewett et al. (2006) have computed the colors of stars in all spectral classes using synthetic photometry (see Bessell 2005 for a description of this technique). To set the zero point, they used the Vega atmospheric model adopted by Bohlin & Gilliland (2004) and assumed that the magnitude of Vega is zero at all wavelengths. This approach has the advantage that corrections can be easily made for adoption of other magnitudes for Vega or an improved atmospheric model for Vega.

Hewett et al. (2006) used an inhomogeneous set of sources in the literature for the spectra of the M, L, and T dwarfs. There is currently a program underway at the IRTF to generate a library of moderate resolution, high signal-to-noise spectra of a large number of stars with reliable spectral types spanning the entire range of known spectral types and luminosity classes. Spectra for M, L, and T dwarfs, included as part of this IRTF spectral library, have already been published (Cushing, Rayner, & Vacca 2005) and spectra for F–T stars are now available on the IRTF Web site <http://irtfweb.ifa.hawaii.edu/~spex/spexlibrary/IRTFlibrary.html>. We are undertaking a project to use these spectra to compute synthetic photometry of the stars in the library to significantly improve on the approach taken by Hewett et al. (2006). The advantages of using this spectral library are (1) it covers the spectral range 0.8–5.0 μm , with resolutions of $R = 2000$ between 0.8–2.5 μm , and $R = 2500$ between 2.5–5.0 μm , and signal-to-noise ratios greater than 100; (2) all of the data were obtained with the same instrument (SpeX, Rayner et al. 2003) and reduced in a consistent manner with the same reduction package (Spextool, Cushing, Vacca, & Rayner 2004); (3) all flux calibrations were achieved using A0V stars, with V mag between 5–8, as spectral standards (Vacca, Cushing, & Rayner 2003); and (4) most of the stars in the spectral library have 2MASS magnitudes available.

Because the IRTF spectral library calibration relies on the model of Vega adopted by Tokunaga & Vacca (2005) and observations of A0V stars, scaled to

their observed V mag and corrected for any possible reddening as estimated from their $B - V$ colors (see Vacca, Cushing, & Rayner 2003), the colors of the program stars as derived from synthetic photometry, should be quite accurate. To determine the systematic uncertainties, we plan to obtain photometry on a set of unreddened A0V stars with the MKO-NIR filters. The average color of these stars would be defined to be zero for all filter combinations. These stars would be then observed in a fashion similar to that used for the IRTF spectral library and compared to the photometric results. Previous comparisons between synthetic photometry and actual measurements indicate that the uncertainties in the colors are at the level of a few percent (Cushing, Vacca, & Rayner 2004).

4. Summary

Infrared photometry is now a mature technique with photometric accuracy comparable to that obtained in the visible wavelength range. However, standardization has not yet been achieved. Combining data from different observations is generally difficult due to the diverse filter sets (often with similar labels) and the numerous photometric systems now in use. We advocate the use of a single filter set, and the widespread use of the MKO-NIR filter set is helping to achieve this. We have summarized some aspects of infrared photometry for the novice to infrared photometry— filter wavelength definitions, the flux density for zero magnitude, atmospheric extinction coefficients, and color corrections to above the atmosphere.

Acknowledgments. A. T. T. acknowledges the support of NASA Cooperative Agreement NCC 5-538. W. D. V. acknowledges partial support from Prof. James R. Graham.

5. APPENDIX A

Web-based photometric information and color transformations can be found at the following locations:

Asiago Database on Photometric Systems:

<http://ulisse.pd.astro.it/Astro/ADPS/Systems/index.html>

Catalog of Infrared Observations:

<http://ircatalog.gsfc.nasa.gov/>

Gemini Near-Infrared Photometric Standard Stars:

<http://www.gemini.edu/sciops/instruments/niri/NIRIPhotStandards.html>

2MASS color transformations:

<http://www.astro.caltech.edu/~jmc/2mass/v3/transformations/>

UKIRT Photometric Calibration:

http://www.jach.hawaii.edu/UKIRT/astronomy/calib/phot_cal/

IRTF photometric information:

http://irtfweb.ifa.hawaii.edu/IRrefdata/ph_catalogs.html;

The MKO-NIR filter profiles can be obtained at:

<http://irtfweb.ifa.hawaii.edu/~nsfcam/hist/filters.2006.html>

References

- Bessell, M. S. 2005, *ARA&A*, 43, 293
- Bessell, M. S., & Brett, J. M. 1988, *PASP*, 100, 1134
- Blackwell, D. E., Leggett, S. K., Petford, A. D., Mountain, C. M., & Selby, M. J. 1983, *MNRAS*, 205, 897
- Bohlin, R. C., & Gilliland, R. L. 2004, *AJ*, 127, 3508
- Carpenter, J. M. 2001, *AJ*, 121, 2851
- Cohen, M., Walker, R. G., Barlow, M. J., & Deacon, J. R. 1992, *AJ*, 104, 1650
- Cohen, M., Megeath, S. T., Hammersley, P. L., Martín-Luis, F., & Stauffer, J. 2003a, *AJ*, 125, 2645
- Cohen, M., Wheaton, W. A., & Megeath, S. T. 2003b, *AJ*, 126, 1090
- Cushing, M. C., Vacca, W. D., & Rayner, J. T. 2004, *PASP*, 116, 362
- Cushing, M. C., Rayner, J. T., & Vacca, W. D. 2005, *ApJ*, 623, 1115
- Elias, J. H., Frogel, J. A., Matthews, K., & Neugebauer, G. 1982, *AJ*, 87, 1029; erratum 1982, *AJ*, 87, 1029
- Fiorucci, M., & Munari, U. 2003, *A&A*, 401, 781
- Fukugita, M., Ichikawa, T., Gunn, J. E., Doi, M., Shimasaku, K., & Schneider, D. P. 1996, *AJ*, 111, 1748
- Glass, I. S. 1999, *Handbook of Infrared Astronomy* (Cambridge: Cambridge University Press)
- Golay, M. 1974, *Introduction to astronomical photometry* (Dordrecht: D. Reidel Publishing Co.)
- Hanner, M. S., Tokunaga, A. T., Veeder, G. J., & Ahearn, M. F. 1984, *AJ*, 89, 162
- Hardie, R. H. 1962, in *Astronomical Techniques*, ed. W. A. Hiltner (Chicago: Univ. of Chicago Press), 178
- Hawarden, T. G., Leggett, S. K., Letawsky, M. B., Ballantyne, D. R., & Casali, M. M. 2001, *MNRAS*, 325, 563
- Henden, A. A., & Kaitchuck, R. H. 1982, *Astronomical Photometry* (New York: Van Nostrand Reinhold)
- Hewett, P. C., Warren, S. J., Leggett, S. K., & Hodgkin, S. T. 2006, *MNRAS*, in press
- Hillenbrand, L. A., Foster, J. B., Persson, S. E., & Matthews, K. 2002, *PASP*, 114, 708
- Hunt, L. K., et al. 1998, *AJ*, 115, 2594
- Johnson, H. L., & Morgan, W. W. 1953, *ApJ*, 117, 313
- Johnson, H. L. 1965, *ApJ*, 141, 923
- Johnson, H. L. 1966, *ARA&A*, 4, 193
- Johnson, H. L., Iriarte, B., Mitchell, R. I., & Wisniewskij, W. Z. 1966, *Comm. of the Lunar and Plan. Lab.*, 4, 99
- Johnson, H. L., MacArthur, J. W., & Mitchell, R. I. 1968, *ApJ*, 152, 465
- Krisciunas, K., et al. 1987, *PASP*, 99, 887
- Leggett, S. K., et al. 2003, *MNRAS*, 345, 144
- Leggett, S. K., et al. 2006, *MNRAS*, preprint
- Low, F. J., & Rieke, G. H. 1974, *The Instrumentation and Techniques of Infrared Photometry*, Vol. 12A (New York: Academic Press)
- Manduca, A., & Bell, R. A. 1979, *PASP*, 91, 848
- Mégessier, C. 1995, *A&A*, 296, 771
- Milone, E. F., & Young, A. T. 2005, *PASP*, 117, 485
- Moro, D., & Munari, U. 2000, *A&AS*, 147, 361
- Peterson, D. M., et al. 2006, *Nat*, 440, 896
- Price, S. D., Paxson, C., Engelke, C., & Murdock, T. L. 2004, *AJ*, 128, 889
- Price, S. D. 2004, *Space Sci.Rev.*, 113, 409
- Rayner, J. T., Toomey, D.W., Onaka, P.M., Denault, A.J., Stahlberger, W.E., Vacca, W.D., Cushing, M.C., & Wang, S. 2003, *PASP*, 115, 362
- Selby, M. J., Mountain, C. M., Blackwell, D. E., Petford, A. D., & Leggett, S. K. 1983, *MNRAS*, 203, 795

- Simons, D. A., & Tokunaga, A. 2002, PASP, 114, 169
Stephens, D. C., & Leggett, S. K. 2004, PASP, 116, 9
Tokunaga, A. T. 2000, in Allen's Astrophysical Quantities, ed. A. N. Cox (4th ed.; New York: Springer), 143
Tokunaga, A. T., Simons, D. A., & Vacca, W. D. 2002, PASP, 114, 180
Tokunaga, A. T., & Vacca, W. D. 2005, PASP, 117, 1459; erratum 2005, PASP, 117, 421
Vacca, W. D., Cushing, M. C., & Rayner, J. T. 2003, PASP, 115, 389
Young, A. T. 1974, in Methods of Experimental Physics: Astrophysics, ed. N. Carleton (New York: Academic Press), 123
Young, A. T., Milone, E. F., & Stagg, C. R. 1994, A&AS, 105, 259

Table 1. Isophotal Wavelengths and Absolute Flux Densities for Vega

Filter	λ_{iso} (μm)	F_{λ} ($\times 10^{-11} \text{W m}^{-2} \mu\text{m}^{-1}$)	ν_{iso} ($\times 10^{14} \text{ Hz}$)	F_{ν} (Jy)
<i>V</i>	0.5450	3680.	5.490	3630
<i>J</i>	1.250	301.	2.394	1560
<i>H</i>	1.644	118.	1.802	1050
<i>K'</i>	2.121	45.7	1.413	686
<i>K_s</i>	2.149	43.5	1.395	670
<i>K</i>	2.198	40.0	1.364	645
<i>L'</i>	3.754	5.31	0.7982	249
<i>M'</i>	4.702	2.22	0.6350	163

Notes: (1) From Tokunaga & Vacca (2005). The uncertainty in the absolute flux density is 1.5%. (2) λ_{iso} and ν_{iso} were computed from using equations (5) and (6). Due to the wide bandpasses of the filters, $\lambda_{\text{iso}} \neq c/\nu_{\text{iso}}$ and $F_{\nu}(\text{Jy}) \neq (\lambda_{\text{iso}}(\mu\text{m}))^2 \times F_{\lambda}(\text{W m}^{-2} \mu\text{m}^{-1})/(3 \times 10^{-12})$.

Table 2. Linear Fit to Calculated Extinction Curve for 2 mm H₂O

Filter	Constant (mag)	Slope (mag/airmass)
<i>J</i>	0.0085 ± 0.0005	0.0153 ± 0.0003
<i>H</i>	0.0091 ± 0.0005	0.0149 ± 0.0003
<i>K</i>	0.0175 ± 0.0008	0.0331 ± 0.0005
<i>K'</i>	0.0731 ± 0.0027	0.0589 ± 0.0015
<i>K_s</i>	0.0442 ± 0.0017	0.0429 ± 0.0010
<i>L'</i>	0.0264 ± 0.0020	0.1039 ± 0.0012
<i>M'</i>	0.1099 ± 0.0049	0.2226 ± 0.0028

Notes: (1) From Tokunaga, Simons, & Vacca (2002). (2) Least-squares linear fitting to the equation (const. + slope*X), where X is the airmass in the range 1.0–3.0.




Figure 1. Comparison of the Johnson (1965) filters (dotted lines) with the MKO-NIR filters (solid lines). Note that the original Johnson infrared filter set did not include the H band. The atmospheric transmission for Mauna Kea with 1 mm precipitable water and an airmass of 1.0 is shown for comparison.




Figure 2. Filter profiles from Stephens & Leggett (2004) illustrating the variations in filter profiles of commonly used photometric systems. The atmospheric transmission is shown by the dotted lines.




Figure 3. MKO-NIR filter extinction curves showing the linear extrapolation to zero airmass (dashed line) for selected filters (adapted from Tokunaga, Simons, & Vacca (2002)). The solid curves show the nonlinear shape of the extinction curve between 0 and 1 airmass; thus extrapolation to zero airmass requires knowledge of the water vapor content of atmosphere above the observatory and stable conditions. Alternatively, narrowband filters such as those proposed by Milone & Young (2005) should be used.